

Factor 13A

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE : IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

DESCRIPTION : Factor XIII in both reduced and non-reduced forms. It does not react with human Factor XIII B-chain or human Factor XII. Factor XIII is a Beta-globulin found in plasma and is composed of two subunits. Factor XIII-A is the catalytic subunit and is a dimer of M.W. Factor XIIIa is a dermal dendrocyte marker and shows variable reaction with these types of tumors.

CATALOG NO :	PL1237	PL1237-R7	7 ML RTU 70 TEST
		PL1237-R1	1 ML RTU 10 TEST
STAINING PATTERN :	Cytoplasmic and Nuclear	PL1237-1	1 ML 1/100 1000 TEST
		PL1237-0,1	0,1 ML 1/100 100 TEST

POSITIVE CONTROL : Recombinant human Factor XIII A-subunit. CAPillary hemangioma, dermatofibroma.

VOLUME : 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide.)

HOST : Mouse

CLONE : AC-1A1

ANTIBODY CONCENTRATION : 200ug / ml

SPECIES REACTIVITY : Human. Others not tested.

EPITOPE : Not determined

MICROBIOLOGICAL STATE : This product is not sterile.

PRETREATMENT : No special pretreatment is required for immuno-histochemistry of formalin-fixed tissues.

PRIMARY ANTIBODY INCUBATION TIME : 30 minutes at Room Temperature

STAINING TIPS : If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY : This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING : Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).