INTENDED USE :   IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

DESCRIPTION :
PMS2 is a gene that encodes for DNA repair proteins involved in mismatch repair. Carriers of the mismatch repair gene mutations have a high lifetime risk of developing Hereditary Non-Polyposis Colon Cancer (HNPCC) and several other cancers including endometrial cancer due to microsatellite instability (MSI) caused by accumulation of DNA replication errors in proliferating cells.

CATALOG NO :   PL2120

PL2120-R7	7 ML RTU 70 TEST
PL2120-R1	1 ML RTU 10 TEST
PL2120-1	1 ML 1/100 1000 TEST
PL2120-0,1	0,1 ML 1/100 100 TEST

STAINING PATTERN :   Nuclear

POSITIVE CONTROL :   Breast, Skin, Cervix, Tonsil, Transitional CellCarcinoma, Colon Carcinoma

VOLUME : 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide.)

HOST :   Mouse

CLONE :   EP51

ANTIBODY CONCENTRATION :   200ug / ml

SPECIES REACTIVITY : Human. Others not tested.

EPITOPE : Not known

MICROBIOLOGICAL STATE :   This product is not sterile.

PRETREATMENT : Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at room temperature for 20 min.

PRIMARY ANTIBODY INCUBATION TIME :   30 minutes at Room Temperature

STAINING TIPS :   If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY : This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING :   Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).