

Calcitonin

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE : IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. It is Rabbit Monoclonal.

DESCRIPTION : Calcitonin is a 32 amino acid peptide which can be demonstrated in C cells of the normal and hyperplastic thyroid. Staining for calcitonin may be used for the identification of a spectrum of C cell proliferative abnormalities ranging from C cell hyperplasia to invasive tumors. Staining for calcitonin in medullary carcinoma of the thyroid produces a fine granular pattern in the cytoplasm. Amyloid deposits within the tumor may also exhibit varying degrees of calcitonin activity.

CATALOG NO : PL9117

PL9117-R7 7 ML RTU 70 TEST

PL9117-R1 1 ML RTU 10 TEST

STAINING PATTERN : Cytoplasmic

PL9117-1 1 ML 1/100 1000 TEST

PL9117-0,1 0,1 ML 1/100 100 TEST

POSITIVE CONTROL : Thyroid or medullary CA

VOLUME : 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide.)

HOST : Rabbit

CLONE : SP17

ANTIBODY CONCENTRATION : Not known

SPECIES REACTIVITY : Human. Others not tested.

EPITOPE : 1-32 aa

MICROBIOLOGICAL STATE : This product is not sterile.

PRETREATMENT : No special pretreatment is required for immuno-histochemistry of formalin-fixed tissues.

PRIMARY ANTIBODY INCUBATION TIME : 30 minutes at Room Temperature

STAINING TIPS : If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY : This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING : Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).