

## CD45 / T200 / LCA

## IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE: IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

## **DESCRIPTION:**

CD45 leucocyte common antigen (LCA)belongs to the family of at least four isoforms of membrane glycoproteins (220, 205, 190, 180kDa) expressed on hematopoietic cell lines but absent on non-hematopoietic cell lines, normal and malignant non-hematopoietic tissues. The intracellular portion of these molecules have protein phosphatase activity and are involved in regulation of transmembrane signals.

CATALOG NO: PL355 PL355 7 ML RTU 70 TEST

PL355-R1 1 ML RTU 10 TEST

**STAINING PATTERN:** Cell Membrane PL355-1 1 ML 1/500 5000 TEST

PL355-0,1 0,1 ML 1/500 500 TEST

**POSITIVE CONTROL**: Tonsil

VOLUME: 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing

stabilizing protein and 0.015mol/L sodium azide.)

**HOST:** Mouse

**CLONE**: PD7/26/16 + 2B11

ANTIBODY CONCENTRATION: 200ug / ml

**SPECIES REACTIVITY:** Human. Others-not tested.

**EPITOPE**: Not determined

MICROBIOLOGICAL STATE: This product is not sterile.

**PRETREATMENT:** No special pretreatment is required for immuno-histochemistry of formalin-fixed tissues.

PRIMARY ANTIBODY INCUBATION TIME: 30 minutes at Room Temperature

STAINING TIPS: If the staining is too light, use lower dilution or longer time. If the staining is

too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY: This product contains sodium azide and is stable for 24 months when stored

at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent

is not stored as recommended, performance must be validated by the user.

**TROUBLESHOOTING:** Please contact Patolab Technical Support by e-mail ( patolab@patolab.com.tr ).



