

## Glycophorin A

### IN VITRO DIAGNOSTIC DATASHEET

**INTENDED USE :** IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy.

**DESCRIPTION :** Glycophorin A is a sialoglycoprotein present on red blood cells (RBC) and erythroid precursor cells. It has been shown that glycophorins acts as the receptor for Sandei virus and parvovirus.

<b>CATALOG NO :</b>	PL1843	PL1843-R7	7 ML RTU 70 TEST
		PL1843-R1	1 ML RTU 10 TEST
<b>STAINING PATTERN :</b>	Cell membrane	PL1843-1	1 ML 1/200 2000 TEST
		PL1843-0,1	0,1 ML 1/200 200 TEST

**POSITIVE CONTROL :** Human erythrocytes in placenta

**VOLUME :** 7 ml Ready to Use ( 7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing stabilizing protein and 0.015mol/L sodium azide. )

**HOST :** Mouse

**CLONE :** JC159

**ANTIBODY CONCENTRATION :** 200ug/ml

**SPECIES REACTIVITY :** Human. Others not tested.

**EPITOPE :** Extracellular domain

**MICROBIOLOGICAL STATE :** This product is not sterile.

**PRETREATMENT :** No special pretreatment is required for the immunohistochemistry of formalin-fixed, paraffin-embedded tissues.

**PRIMARY ANTIBODY INCUBATION TIME :** 30 minutes at Room Temperature

**STAINING TIPS :** If the staining is too light, use lower dilution or longer time. If the staining is too strong, check pretreatment, use higher dilution or shorter time.

**STORAGE AND STABILITY :** This product contains sodium azide and is stable for 24 months when stored at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent is not stored as recommended, performance must be validated by the user.

**TROUBLESHOOTING :** Please contact Patolab Technical Support by e-mail ( patolab@patolab.com.tr ).