

Keratin 5

IN VITRO DIAGNOSTIC DATASHEET

INTENDED USE: IN VITRO DIAGNOSTIC USE

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. Rabbit Monoclonal

DESCRIPTION:

This Antibody recognizes a protein of 58kDa, which is identified as Cytokeratin 5 (KRT5). This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member Keratin14.

CATALOG NO: PL1896 PL1896-R7 7 ML RTU 70 TEST

PL1896-R1 1 ML RTU 10 TEST

STAINING PATTERN: Cytoplasmic PL1896-1 1 ML 1/100 1000 TEST

PL1896-0,1 0,1 ML 1/100 100 TEST

POSITIVE CONTROL: Prostate

VOLUME: 7 ml Ready to Use (7 ml of antibody prediluted in 0.05mol/L Tris-HCl, pH 7.6 containing

stabilizing protein and 0.015mol/L sodium azide.)

HOST: Rabbit

CLONE: ZR280

ANTIBODY CONCENTRATION: Not known

SPECIES REACTIVITY: Human. Others not-tested

EPITOPE: Not determined

MICROBIOLOGICAL STATE: This product is not sterile.

PRETREATMENT: Staining of formalin-fixed tissue sections requires treating the tissue sections in boiling 1mM EDTA, pH 8.0, for 10-20 min followed by cooling at RT for 20 min.

PRIMARY ANTIBODY INCUBATION TIME: 60 minutes at Room Temperature

STAINING TIPS: If the staining is too light, use lower dilution or longer time. If the staining is

too strong, check pretreatment, use higher dilution or shorter time.

STORAGE AND STABILITY: This product contains sodium azide and is stable for 24 months when stored

at 2-8°C. Do not use after expiration date indicated on label of the product. If reagent

is not stored as recommended, performance must be validated by the user.

TROUBLESHOOTING: Please contact Patolab Technical Support by e-mail (patolab@patolab.com.tr).



